

Educando para a paz

Autores Autores Autor (es) USF Lucas de Francisco Carvalho Ana Devvis Santos Araújo Jesuíno Samanta Romanin Zuchetto Ana Carolina Zuanazzi Ana Carolina Zuanazzi Autores Internacionais Programa de Pós Graduação Stricto Sensu em Psicologia DOI 10.15448/1980-8623.2021.4.36442 Assunto (palavras chaves) personality disorders, differential diagnosis, psychopathy, psychological assessment Idioma Português Fonte Título do periódico: Psico ISSN: 1980-8623 Volume/Número/Paginação/Ano: 52/e36442/2021 Data da publicação 19-1-2022 Formato da produção Digital Resumo We aimed to develop a version of the Dimensional Clinical Personality Inventory 2 (IDCP-2) according to the Hierarchical Taxonomy of Psychopathology (HiTOP) for the assessment of antisocial personality disorder traits (i.e., IDCP Antisocial Personality Disorder Scale; IDCP-ASPD), as well as verify its psychometric properties. We developed new factors to cover ASPD traits, and collect data with 206 adults from the community (Mage = 31.3; 77.8% women). Participants completed the IDCP-ASPD, factors from IDCP-2, and Personality Inventory for DSM-5 (PID-5). Exploratory structural equation modeling (E-SEM) suggested a 3 factors solution, grouping the 14 factors composing the IDCP-ASPD. Raive as found for the usability of the developed scale for ASPD traits measures corroborated expectations. The bootstrap two-sample t-test comparing non-clinical and psychiatric	Tipo	Periódico
Autor (es) USF Lucas de Francisco Carvalho Ana Deyvis Santos Araújo Jesuíno Samanta Romanin Zuchetto Autores Internacionais Programa de Pós Graduação Stricto Sensu em Psicologia DOI 10.15448/1980-8623.2021.4.36442 Assunto (palavras chaves) personality disorders, differential diagnosis, psychopathy, psychological assessment Idioma Português Fonte Título do periódico: Psico ISSN: 1980-8623 Volume/Número/Paginação/Ano: 52/e36442/2021 Data da publicação 19-1-2022 Formato da produção Digital Resumo We aimed to develop a version of the Dimensional Clinical Personality Inventory 2 (IDCP-2) according to the Hierarchical Taxonomy of Psychopathology (HITOP) for the assessment of antisocial personality disorders to cover ASPD traits, and collect data with 206 adults from the community (Mage = 31.3; 77.8% women). Participants completed the IDCP-ASPD, factors from IDCP-2, and Personality Inventory for DSM-5 (PID-5). Exploratory structural equation modeling (E-SEM) suggested a 3 factors solution, grouping the 14 factors composing the IDCP-ASPD and external measures corroborated expectations. The bootstrap two-sample t-test comparing non-clinical and psychiatric groups suggested good discrimination capacity of the IDCP-ASPD. Favorable evidence was found for the usability of the developed scale for ASPD traits measurement, although future studies must replicate the findings in samples composed by ASPD patients.	Título	
Ana Deyvis Santos Araújo Jesuíno Samanta Romanin Zuchetto Ana Carolina ZuanazziAutores InternacionaisPrograma/Curso (s)Programa de Pós Graduação Stricto Sensu em PsicologiaDOI10.15448/1980-8623.2021.4.36442Assunto (palavras chaves)personality disorders, differential diagnosis, psychopathy, psychological assessmentIdiomaPortuguêsFonteTítulo do periódico: Psico ISSN: 1980-8623 Volume/Número/Paginação/Ano: 52/e36442/2021Data da publicação19-1-2022Formato da produçãoDigital We aimed to develop a version of the Dimensional Clinical Personality Inventory 2 (IDCP-2) according to the Hierarchical Taxonomy of Psychopathology (HITOP) for the assessment of antisocial personality disorder traits (.e., IDCP Antisocial Personality Disorder Scale; IDCP-ASPD, as well as verify its psychometric properties. We developed new factors to cover ASPD traits, and collect data with 206 adults from the community (Mage = 31.3; 77.8% women). Participants completed the IDCP-ASPD, factors from IDCP-2, and Personality Inventory for DSM-5 (PID-5). Exploratory structural equation modeling (E-SEM) suggested a 3 factors solution, grouping the 14 factors composing the IDCP- ASPD. Reliability indicators were good. Correlations between IDCP-ASPD and external measures corroborated expectations. The bootstrap two-sample t-test comparing non-clinical and psychiatric groups suggested good discrimination capacity of the IDCP-ASPD. Favorable evidence was found for the usability of the developed scale for ASPD traits measurement, although future studies must replicate the findings in samples composed by ASPD patients.	Autores	
Programa/Curso (s) Programa de Pós Graduação Stricto Sensu em Psicologia DOI 10.15448/1980-8623.2021.4.36442 Assunto (palavras personality disorders, differential diagnosis, psychopathy, psychological assessment Idioma Português Fonte Título do periódico: Psico ISSN: 1980-8623 Volume/Número/Paginação/Ano: 52/e36442/2021 Data da publicação 19-1-2022 Formato da produção Digital Resumo We aimed to develop a version of the Dimensional Clinical Personality Inventory 2 (IDCP-2) according to the Hierarchical Taxonomy of Psychopathology (HiTOP) for the assessment of antisocial personality disorder traits (i.e., IDCP Antisocial Personality Disorder Scale; IDCP-ASPD), as well as verify its psychometric properties. We developed new factors to cover ASPD traits, and collect data with 206 adults from the community (Mage = 31.3; 77.8% women). Participants completed the IDCP-ASPD, factors from IDCP-2, and Personality Inventory for DSM-5 (PID-5). Exploratory structural equation modeling (E-SEM) suggested a 3 factors solution, grouping the 14 factors composing the IDCP-ASPD. Reliability indicators were good. Correlations between IDCP-ASPD and external measures corroborated expectations. The bootstrap two-sample t-test comparing non-clinical and psychiatric groups suggested good discimination capacity of the IDCP-ASPD. Favorable evidence was found for the usability of the developed scale for ASPD. Traits measurement, although future studies must replicate the findings in samples composed by ASPD patients.	Autor (es) USF	Ana Deyvis Santos Araújo Jesuíno Samanta Romanin Zuchetto
DOI 10.15448/1980-8623.2021.4.36442 Assunto (palavras chaves) personality disorders, differential diagnosis, psychopathy, psychological assessment Idioma Português Fonte Título do periódico: Psico ISSN: 1980-8623 Volume/Número/Paginação/Ano: 52/e36442/2021 Data da publicação 19-1-2022 Formato da produção Digital We aimed to develop a version of the Dimensional Clinical Personality Inventory 2 (IDCP-2) according to the Hierarchical Taxonomy of Psychopathology (HiTOP) for the assessment of antisocial personality disorder traits (i.e., IDCP Antisocial Personality Disorder Scale; IDCP-ASPD), as well as verify its psychometric properties. We developed new factors to cover ASPD traits, and collect data with 206 adults from the community (Mage = 31.3; 77.8% women). Participants completed the IDCP-ASPD, factors from IDCP-2, and Personality Inventory for DSM-5 (PID-5). Exploratory structural equation modeling (E-SEM) suggested a 3 factors solution, grouping the 14 factors composing the IDCP-ASPD. Reliability indicators were good. Correlations between IDCP-ASPD and external measures corroborated expectations. The bootstrap two-sample t-test comparing non-clinical and psychiatric groups suggested good discrimination capacity of the IDCP-ASPD. Favorable evidence was found for the usability of the developed scale for ASPD. Traits measurement, although future studies must replicate the findings in samples composed by ASPD patients.	Autores Internacionais	
Assunto (palavras chaves)personality disorders, differential diagnosis, psychopathy, psychological assessment assessmentIdiomaPortuguêsFonteTítulo do periódico: Psico 	Programa/Curso (s)	Programa de Pós Graduação Stricto Sensu em Psicologia
chaves) assessment Idioma Português Fonte Título do periódico: Psico ISSN: 1980-8623 Volume/Número/Paginação/Ano: 52/e36442/2021 Data da publicação 19-1-2022 Formato da produção Digital Resumo We aimed to develop a version of the Dimensional Clinical Personality Inventory 2 (IDCP-2) according to the Hierarchical Taxonomy of Psychopathology (HiTOP) for the assessment of antisocial personality disorder traits (i.e., IDCP Antisocial Personality Disorder Scale; IDCP-ASPD), as well as verify its psychometric properties. We developed new factors to cover ASPD traits, and collect data with 206 adults from the community (Mage = 31.3; 77.8% women). Participants completed the IDCP-ASPD, factors from IDCP-2, and Personality Inventory for DSM-5 (PID-5). Exploratory structural equation modeling (E-SEM) suggested a 3 factors solution, grouping the 14 factors composing the IDCP- ASPD. Reliability indicators were good. Correlations between IDCP-ASPD and external measures corroborated expectations. The bootstrap two-sample t-test comparing non-clinical and psychiatric groups suggested good discrimination capacity of the IDCP-ASPD. Favorable evidence was found for the usability of the developed scale for ASPD traits measurement, although future studies must replicate the findings in samples composed by ASPD patients.	DOI	10.15448/1980-8623.2021.4.36442
FonteTítulo do periódico: Psico ISSN: 1980-8623 Volume/Número/Paginação/Ano: 52/e36442/2021Data da publicação19-1-2022Formato da produçãoDigitalResumoWe aimed to develop a version of the Dimensional Clinical Personality Inventory 2 (IDCP-2) according to the Hierarchical Taxonomy of Psychopathology (HiTOP) for the assessment of antisocial personality disorder traits (i.e., IDCP Antisocial Personality Disorder Scale; IDCP-ASPD), as well as verify its psychometric properties. We developed new factors to cover ASPD traits, and collect data with 206 adults from the community (Mage = 31.3; 77.8% women). Participants completed the IDCP-ASPD, factors from IDCP-2, and Personality Inventory for DSM-5 (PID-5). Exploratory structural equation modeling (E-SEM) suggested a 3 factors solution, grouping the 14 factors composing the IDCP- ASPD. Reliability indicators were good. Correlations between IDCP-ASPD and external measures corroborated expectations. The bootstrap two-sample t-test comparing non-clinical and psychiatric groups suggested good discrimination capacity of the IDCP-ASPD. Favorable evidence was found for the usability of the developed scale for ASPD traits measurement, although future studies must replicate the findings in samples composed by ASPD patients.	Assunto (palavras chaves)	
ISSN: 1980-8623 Volume/Número/Paginação/Ano: 52/e36442/2021Data da publicação19-1-2022Formato da produçãoDigitalResumoWe aimed to develop a version of the Dimensional Clinical Personality Inventory 2 (IDCP-2) according to the Hierarchical Taxonomy of Psychopathology (HiTOP) for the assessment of antisocial personality disorder traits (i.e., IDCP 	Idioma	
Formato da produçãoDigitalResumoWe aimed to develop a version of the Dimensional Clinical Personality Inventory 2 (IDCP-2) according to the Hierarchical Taxonomy of Psychopathology (HiTOP) for the assessment of antisocial personality disorder traits (i.e., IDCP Antisocial Personality Disorder Scale; IDCP-ASPD), as well as verify its psychometric properties. We developed new factors to cover ASPD traits, and collect data with 206 adults from the community (Mage = 31.3; 77.8% women). Participants completed the IDCP-ASPD, factors from IDCP-2, and Personality Inventory for DSM-5 (PID-5). Exploratory structural equation modeling (E-SEM) suggested a 3 factors solution, grouping the 14 factors composing the IDCP- ASPD. Reliability indicators were good. Correlations between IDCP-ASPD and external measures corroborated expectations. The bootstrap two-sample t-test comparing non-clinical and psychiatric groups suggested good discrimination capacity of the IDCP-ASPD. Favorable evidence was found for the usability of the developed scale for ASPD traits measurement, although future studies must replicate the findings in samples composed by ASPD patients.	Fonte	ISSN: 1980-8623
Resumo We aimed to develop a version of the Dimensional Clinical Personality Inventory 2 (IDCP-2) according to the Hierarchical Taxonomy of Psychopathology (HiTOP) for the assessment of antisocial personality disorder traits (i.e., IDCP Antisocial Personality Disorder Scale; IDCP-ASPD), as well as verify its psychometric properties. We developed new factors to cover ASPD traits, and collect data with 206 adults from the community (Mage = 31.3; 77.8% women). Participants completed the IDCP-ASPD, factors from IDCP-2, and Personality Inventory for DSM-5 (PID-5). Exploratory structural equation modeling (E-SEM) suggested a 3 factors solution, grouping the 14 factors composing the IDCP-ASPD. Reliability indicators were good. Correlations between IDCP-ASPD and external measures corroborated expectations. The bootstrap two-sample t-test comparing non-clinical and psychiatric groups suggested good discrimination capacity of the IDCP-ASPD. Favorable evidence was found for the usability of the developed scale for ASPD traits measurement, although future studies must replicate the findings in samples composed by ASPD patients.	Data da publicação	19-1-2022
Resumo We aimed to develop a version of the Dimensional Clinical Personality Inventory 2 (IDCP-2) according to the Hierarchical Taxonomy of Psychopathology (HiTOP) for the assessment of antisocial personality disorder traits (i.e., IDCP Antisocial Personality Disorder Scale; IDCP-ASPD), as well as verify its psychometric properties. We developed new factors to cover ASPD traits, and collect data with 206 adults from the community (Mage = 31.3; 77.8% women). Participants completed the IDCP-ASPD, factors from IDCP-2, and Personality Inventory for DSM-5 (PID-5). Exploratory structural equation modeling (E-SEM) suggested a 3 factors solution, grouping the 14 factors composing the IDCP-ASPD. Reliability indicators were good. Correlations between IDCP-ASPD and external measures corroborated expectations. The bootstrap two-sample t-test comparing non-clinical and psychiatric groups suggested good discrimination capacity of the IDCP-ASPD. Favorable evidence was found for the usability of the developed scale for ASPD traits measurement, although future studies must replicate the findings in samples composed by ASPD patients.	Formato da produção	Digital
	Resumo	We aimed to develop a version of the Dimensional Clinical Personality Inventory 2 (IDCP-2) according to the Hierarchical Taxonomy of Psychopathology (HiTOP) for the assessment of antisocial personality disorder traits (i.e., IDCP Antisocial Personality Disorder Scale; IDCP-ASPD), as well as verify its psychometric properties. We developed new factors to cover ASPD traits, and collect data with 206 adults from the community (Mage = 31.3; 77.8% women). Participants completed the IDCP-ASPD, factors from IDCP-2, and Personality Inventory for DSM-5 (PID-5). Exploratory structural equation modeling (E-SEM) suggested a 3 factors solution, grouping the 14 factors composing the IDCP-ASPD. Reliability indicators were good. Correlations between IDCP-ASPD and external measures corroborated expectations. The bootstrap two-sample t-test comparing non-clinical and psychiatric groups suggested good discrimination capacity of the IDCP-ASPD. Favorable evidence was found for the usability of the developed scale for ASPD traits measurement, although future studies must
	Fomento	<u> </u>



+ +

+

+